

Clifford geometric parameterization of inequivalent vacua

Bertfried Fauser
Universität Konstanz
Fakultät für Physik
Fach M 678

78457 Konstanz **Germany**

Electronic mail: Bertfried.Fauser@uni-konstanz.de

Feb 14, 2001

Abstract

We propose a geometric method to parameterize inequivalent vacua by dynamical data. Introducing quantum Clifford algebras with arbitrary bilinear forms we distinguish isomorphic algebras –as Clifford algebras– by different filtrations resp. induced gradings. The idea of a vacuum is introduced as the unique algebraic projection on the base field embedded in the Clifford algebra, which is however equivalent to the term vacuum in axiomatic quantum field theory and the GNS construction in C^* -algebras. This approach is shown to be equivalent to the usual picture which fixes one product but employs a variety of GNS states. The most striking novelty of the geometric approach is the fact that dynamical data fix uniquely the vacuum and that positivity is not required. The usual concept of a statistical quantum state can be generalized to geometric meaningful but non-statistical, non-definite, situations. Furthermore, an algebraization of states takes place. An application to physics is provided by an $U(2)$ -symmetry producing a gap-equation which governs a phase transition. The parameterization of all vacua is explicitly calculated from propagator matrix elements. A discussion of the relation to BCS theory and Bogoliubov-Valatin transformations is given.

PACS: 03.70; 11.10; 64.10

MSC2000: 15A66; 81S05; 81R99

Keywords: Clifford algebras, CAR algebras, inequivalent vacua, state space, phase transitions, gap-equation, indefinite states, geometric interpretation of quantum mechanics, multivectors, $\mathbb{Z}_n$ -grading, GNS states, BCS theory, Bogoliubov-Valatin transformation

1 Introduction

1.1 Prologue

The core material of this paper, mainly done in 1997, originated from an analysis of the C^* -algebraic considerations of Kerschner [31, 32] and had a major impact on the progress of a whole set of further work, supporting the need of an exposition of the method itself. Kerschner reflected the need of a more involved concept of (ground) state or vacuum in quantum field theory. Due to Haag's theorem an interacting quantum field theory (QFT) cannot be treated in Fock space. Since Kerschner and the group of Prof. Stumpf were involved in composite calculations, see [42], this problem was serious. However, since regularization issues made it necessary to perform calculations in an indefinite state space C^* -algebraic methods are not appropriate because they are tied to positive definiteness which causes problems. It proved to be a very fruitful concept to turn this considerations into a Clifford algebraic picture. A reformulation of QFT was done in [13]. It was necessary to enlarge the concept of Clifford algebra to arbitrary bilinear forms which are not necessary symmetric, see e.g. [14, 1, 37]. Later on, Dirac theory [16], composite calculations in quantum electro dynamics [15] and normal-ordering in quantum field theory [12, 24] have been treated. Surprisingly several singularities which usually occur during re-ordering in QFT vanished due to the correct algebraic treatment. A general method to deal with QFT in Clifford algebraic terms was given in [17]. The method allowed to generalize some concepts of QFT. Especially it was shown [18, 19, 3, 21] that q -symmetry can be treated within this formalism. A plain approach to the notions and peculiarities of quantum Clifford algebras was given in [20]. Some further achievements will be discussed in the epilogue. All this applications have its foundation in the method which is presented in the present work. A low dimensional model which shows all peculiarities of QFT while being still fully accessible to calculations seemed to be the perfect playground to demonstrate the method since a comparison to C^* -algebraic methods is also easily possible.

We want to add a few remarks on notational and conceptual issues about the concept of **vacuum** used here:

- The term vacuum could be called effective or generalized vacuum to express some differences with conventional settings. But from a physicists point of view it is simply the unique state which is used to generate a state space and calculate expectation values. This is the physical vacuum of the theory.
- If the vacuum is seen as a ground state or a cyclic lowest weight vector one assumes a Hamiltonian and some annihilation operators which annihilate this state. Since we do not choose a particular Hamiltonian but deal with a whole class of dynamics invariant under a group action, e.g. $U(2)$, one cannot characterize vacua in this way. Moreover, in non-Fock situations one cannot expect to find annihilators in terms of the field operators, see below.
- A basic difference between the C^* -algebraic approach and our method is that in C^* -theory one deals with a state *space*, i.e. a convex set of positive linear functionals. An operator equation can in principle be evaluated with any state from this set. But in our approach the method yields an *unique* state and no freedom is possible. This is a major achievement since a unique representation is

chosen from a possibly infinite set. The theory is thus more predictive. A unique state fixes properties of a quantum field theory as e.g. symmetry breaking or forming of a condensate. This connection is established via the dynamics of the theory which allows to speak about a dynamically chosen representation.

- The last point has to be supplemented by the observation that our vacuum state cannot be addressed as a thermalized state as it is done in thermo field dynamics.

1.2 Motivation

Quantum physics arose from the necessity to generalize coordinates and coordinate functions to operators and operator valued functions. The historical development of quantum mechanics led to the concept of a Hilbert space, a linear space with a positive definite inner product, or scalar product, usually over the complex number field [36, 10]. The so-called observables are then no longer differentiable functions on a phase-space, but bounded, hermitean operators on an appropriate Hilbert space [36]. We want to emphasize here, that operators are not observables in their own right. From an empirical point of view, let's assume a measurement yields a real number. A measurable quantity is thus a basis dependent expectation value e.g. a matrix element or an invariant object which is not basis dependent, e.g. an eigenvalue. Only in the positive definite case of a Hilbert space, are we able to reconstruct the operators up to isomorphisms, hence their invariants, uniquely from such data; finite dimension or separability is also assumed. This *a posteriori* identification allows us to address the bounded operators on a Hilbert space as observables. We will be concerned here with a geometric relation between operators and their invariants.

In quantum field theory, however, one fails to find such a direct connection. The canonical (anti) commutation relations CCR (CAR) of a continuum of variables –the space-time continuum has to be considered as an index– do possess infinitely many representations. One does therefore lose the information about a particular representation when passing from expectation values to operators and possibly an operator dynamics. On the other hand, the freedom in choosing a representation can be used to treat fields at finite temperature or within different phases and provides therefore an ideal tool in modelling physical systems. Furthermore, it is known from Haag's theorem, that interacting QFT's cannot be described correctly within Fock space [27, 5]. It is thus of utmost importance to have a *constructive* way to handle non-Fock situations. Troubled with the requirement of positivity, in the relativistic setting, these situations are usually handled with complicated inductive limits in a C^* -algebraic approach [28]. Our geometric approach will allow such intriguing situations in the finite dimensional case also, which might be used to circumvent convergence problems.

The need of positivity and definiteness results from another point also. Quantum mechanics was developed within a background of positivism and it was welcomed by its founders to introduce a statistical and *a priori* unquestionable element into the theory –the chance. Especially Bohr and the so-called Copenhagen interpretation developed by Born took this route. Nevertheless, already in the thirties and mostly connected with the appearance of the electron spin, geometrical concepts were investigated [26]. We will return to geometric conceptions which include, however, the QM case as a

subset.

In this paper we give a geometric interpretation of vacuum states and their relation to symmetry and the dynamics under consideration within a Clifford algebraic framework. However, much effort is spent to show that in the positive case of QM our approach reproduces the common results, even if it is in some sense more general. This method will include a geometric interpretation of “quantization” also. We will concentrate on finite dimensional examples i.e. fermions for reasons of clarity and to avoid convergence problems, which do not belong to the geometric picture. However, the method can be directly applied to quantum field theory [15, 12, 13]. Furthermore, one should note, that in our framework all the interesting phenomena as phase transitions etc. can be handled with finitely many particles.

The striking advantage of the geometric picture is that it is neither connected to positivity nor to definiteness. If we artificially fix the situation in the positive definite case, which can be handled in the usual way, we are ready to change to the geometric picture and afterwards extend the theory to formerly meaningless situations. Since in quantum field theory ghost states –of negative and zero norm– are presently used e.g. in QCD, we expect a clarification of these situations by our approach.

An algebraic definition of the term vacuum is needed, which includes conventional vacua, but is accessible also to the general algebraic picture. The most important feature is invariance under a symmetry group, which is in axiomatic QFT the Lorentz group. We will use for simplicity the nonrelativistic case and employ the $U(2)$. The construction of states will be done by an analogy to C^* -algebraic GNS states [35], which however are generalized to indefinite cases by Clifford geometric methods. As a major outcome, which distinguishes our method from any other current approach, the Clifford algebra will provide an unique vacuum for every dynamical situation, enabling us to connect propagator data to vacuum states.

Since we are presenting a geometric approach to the concept of states and furthermore their algebraic treatment, we have to compare conventional and new calculations. Therefore, in section 2 an explicit treatment of the one dimensional extended Fermi oscillator is given, first to fix the notations and second for didactical reasons, since this is the model where old and new approaches share most of their features.

Section 3 is devoted to the development of the geometric theory in the finite dimensional case, which however can easily be used in QFT by analogy. A resume of quantum Clifford algebras with arbitrary bilinear forms is given in section 3.1. Sections 3.2 and 3.3 treat once more the Fermi oscillator, simply by translating the results from 2 and only thereafter turning to a radical geometric point of view. It seemed to be necessary to perform this transition in two steps, to be as clear as possible. A redundancy occurs only in the results, but not in the novel points of view. Section 3.4 provides a treatment of the Fermi oscillator with two degrees of freedom, first giving the results of ECAR calculations and then a complete discussion within the geometric picture. The four main results given there are a description of inequivalent vacua in a parametric and constructive form, the explicit calculation of the functional dependency of these parameters from the matrix elements of the propagator, a ‘gap-equation’ which distinguishes different phases, and a complete classification of all vacuum states by geometrical data.

The 4th section gives the connection of our approach to BCS theory and Bogoliubov-

Valatin transformations as a relation to results from literature. Furthermore, it speculates, how this approach may be related to the Higgs mechanism and ghost fields.

2 The extended Fermi oscillator

Let us consider the ordinary Fermi oscillator with one degree of freedom. This very trivial system serves us finding the relevant relations by explicit calculations.

The CAR algebra of one degree of freedom is generated by $\mathbb{1}$, and $a, a^\dagger$, where the relations

$$\begin{aligned} aa^\dagger + a^\dagger a &= \mathbb{1} \\ aa &= 0 = a^\dagger a^\dagger \end{aligned} \quad (1)$$

hold. We will deal with the real field for simplicity, but a complexification could easily be performed by $CAR_{\mathbb{C}} = \mathbb{C} \otimes CAR_{\mathbb{R}}$, where bilinear forms have to be changed into sesquilinear ones.

We find an algebraic basis of the algebra which spans a (real) linear space $FCAR$ of dimension $2^n = 4$ – using the forgetful functor F . Generally we will write FA for the linear space underlying the algebra A –

$$\{e_i\}_{i=1}^4 = \{\mathbb{1}, aa^\dagger, a, a^\dagger\}. \quad (2)$$

A state ω on this algebra is an element of the linear forms $\text{lin}[FCAR, \mathbb{R}] \simeq FCAR^*$, where $*$ does indicate the dual with respect to a canonical pairing. It is usually supposed to be linear, positive and normed. Since $FCAR$ and $FCAR^*$ are isomorphic as linear spaces, any dual element can be parameterized by its action on the basis elements in $FCAR$. Introducing an *independent* “particle number” operator or even better a grading by

$$\begin{aligned} [Q, a] &= -a \\ [Q, a^\dagger] &= +a^\dagger \\ Q &\notin \text{Alg}(a, a^\dagger), \end{aligned} \quad (3)$$

we define the Extended CAR (ECAR) algebra [34]. Regarding Q as an independent quantity, allows us to have circumvented von Neumann’s uniqueness theorem and provides inequivalent vacua in the finite dimensional case also. This is also the place where deformations, especially the q -business, enters the setting [19, 18].

A CAR algebra subjected to the action of a group G here $SO(2)$ resp. $U(1)$ generated by a set of generators e.g. Q , is called a CAR-dynamical system.

A *vacuum state* ω is defined to be a state invariant under the action of the group G here a $U(1)$ introduced by Q . The invariance condition reads

$$\omega(A) = \omega(QA) \quad (= \omega(AQ)). \quad (4)$$

This is exactly the definition of a vacuum due to axiomatic quantum field theory [41]. An emphasis is laid on symmetry principles and not on step operators –creation and

annihilation operators— or ground states which are annihilated by the later. See also the discussion of Bogoliubov-Valatin transformations and of the BCS vacuum below.

In the case of axiomatic QFT [44], the desired group would be the Lorentz group, however, we deal with $U(1)$ and $U(2)$ for simplicity. The $U(2)$ example is already of physical interest and models superconductivity. Furthermore, if a particle interpretation is desired, the vacuum state has to be a highest (lowest) weight state of a specific representation of the group acting in our system. This is in fact the GNS state $|\Omega_k\rangle \cong |0\rangle_{GNS}$ of a *specific* representation π . However, we also deal with indefinite situations. A lowest weight state is called *ground state* for physical systems and will be denoted as $|0\rangle$. It obeys the Fock-like condition $X^- |0\rangle = 0$ for some ladder operator X^- . Furthermore, from (4) we can conclude that a vacuum state has no nontrivial quantum number w.r.t. the chosen group. State and group action define an ECAR dynamical system, if additionally a Hamiltonian element is specified. However we will see that it is sufficient to specify the propagator of the theory *as a parameter* which fully specifies the vacuum properties. This relation leading to multiple Hamiltonians bearing the same vacuum structure is not injective. In the sequel, we deal exclusively with vacuum states denoting them loosely as $S^0(ECAR) = \{\omega_k\}$, with k an index set.

This means that we seek for vacuum states, which are defined to posses no nonzero G quantum numbers. Together with the invariance condition (4) which leads to $\omega(AQ) = 0, \forall A \in ECAR$, we have to fulfil in our case:

$$\begin{aligned} i) \quad & \omega(\mathbb{1}) = 1 && \text{normalization} \\ ii) \quad & \omega(a) = 0 = \omega(a^\dagger) && \text{due to the grading} \\ iii) \quad & \omega(aa^\dagger) = \nu && \end{aligned} \tag{5}$$

where ν is a real parameter. The positivity requirement restricts ν by $\nu > 0$ and $\omega(a^\dagger a) = \omega(\mathbb{1} - aa^\dagger) = 1 - \nu > 0$ which results in $1 > \nu > 0$. Hence we can parameterize all linear, normed, positive vacuum states $S^0(ECAR)$ by a single real quantity $\nu \in]0, 1[$. We write ω_ν for such *vacuum states* in $S^0(ECAR)$.

We can now calculate ν -dependent pseudo representation matrices by using ω_ν as linear form and the $\{e_i\}$ basis. With $A \in ECAR$ we have

$$[\mu_{ij}^\nu(A)] = [\omega_\nu(e_i^\dagger A e_j)] \tag{6}$$

e.g.

$$[\eta_{ij}^\nu] = [\mu_{ij}^\nu(\mathbb{1})] = \begin{vmatrix} 1 & \nu & 0 & 0 \\ \nu & \nu & 0 & 0 \\ 0 & 0 & 1 - \nu & 0 \\ 0 & 0 & 0 & \nu \end{vmatrix} \quad [\mu_{ij}^\nu(a)] = \begin{vmatrix} 0 & 0 & 0 & \nu \\ 0 & 0 & 0 & \nu \\ 1 - \nu & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{vmatrix} \tag{7}$$

This is not a homomorphism and the algebra product is *not* mapped onto the matrix

product, so μ is not a representation, which can be seen from

$$\begin{aligned}
\omega_\nu(e_i^\dagger A B e_j) &= \omega_\nu(e_i^\dagger A e_k) b_{jk} \\
&= \omega_\nu(e_i^\dagger A e_r) \eta_{rl}^{-1} \eta_{lk} b_{jk} \\
&= \omega_\nu(e_i^\dagger A e_r) \eta_{rl}^{-1} b_{jk} \omega_\nu(e_l^\dagger e_k) \\
&= \omega_\nu(e_i^\dagger A e_r) \eta_{rl}^{-1} \omega(e_l^\dagger B e_j)
\end{aligned} \tag{8}$$

or equivalently

$$\mu(AB) = \mu(A) \circ \eta^{-1} \circ \mu(B). \tag{9}$$

As long as η is nonsingular and positive, which is not the case in Fock or dual Fock space or for values of $\nu \notin]0, 1[$, see below, one could proceed to a $*$ -representation π by setting

$$\pi^\nu := \eta^{-1/2} \circ \mu^\nu \circ \eta^{-1/2}, \quad \nu \in]0, 1[. \tag{10}$$

The two extremal and singular cases $\nu = 0, 1$ reduce the situation to usual Fock F , $\pi^1(a) \mid 0 >_F = 0$ and dual Fock F^* , $\pi^0(a^\dagger) \mid 0 >_{F^*} = 0$ space after factoring out a null-space. A connection between them is achieved by renaming $a, a^\dagger$ to $a^\dagger, a$. In general, if $\nu \neq 0$ or 1 , the representations are no longer reducible and remain to be four dimensional. However, a decomposition into Fock and dual-Fock spaces can be performed using mixed states. Note once more, that $Q \neq Q(a, a^\dagger)$ is an independent quantity. However, from the group action (3) we see that $(ad_Q)^2 = Id$ on $V = \langle a, a^\dagger \rangle$ and the basis set is only $\{e_i\}_{i=1}^4 \oplus \{e_i\}_{i=1}^4 Q$.

At the end of this section, we give a further possibility to describe the ν -dependent vacua, which is needed later. From a decomposition of unity into primitive projectors we have

$$\mathbb{1} = P_F + P_{F^*} \tag{11}$$

and the decomposition

$$\begin{aligned}
\omega_\nu(A^\dagger B) &= \omega_\nu(A^\dagger [P_F + P_{F^*}] B) \\
&= \omega_\nu(A^\dagger P_F B) + \omega_\nu(A^\dagger P_{F^*} B) \\
&=: \nu \omega_F(A^\dagger B) + (1 - \nu) \omega_{F^*}(A^\dagger B),
\end{aligned} \tag{12}$$

where ω_F and ω_{F^*} are the extremal Fock and dual Fock states. If one likes to proceed to a diagonal decomposition it is convenient furthermore to introduce GNS ground states $|\Omega_\nu\rangle$. This results up to a relative phase in

$$\begin{aligned}
\omega_\nu(A) &= \langle \Omega_\nu \mid \pi_\nu(A) \Omega_\nu \rangle \\
&= \nu \langle \Omega_F \mid \pi_F(A) \Omega_F \rangle + (1 - \nu) \langle \Omega_{F^*} \mid \pi_{F^*}(A) \Omega_{F^*} \rangle
\end{aligned} \tag{13}$$

$$\pi_\nu(A) = \pi_F(A) \oplus \pi_{F^*}(A) \tag{14}$$

$$|\Omega_\nu\rangle = \sqrt{\nu} |\Omega_F\rangle \oplus \sqrt{1 - \nu} |\Omega_{F^*}\rangle. \tag{15}$$

Since Fock and dual Fock situations are connected by a simple relabelling of generators, the distinction seems to be a commonplace. But if more generators are involved other representations come into play, furthermore the concrete expression of Q depends already on this choice. However, one should be careful in transforming generators, since mathematical features such as simplicity of the generated multiplicative groups may depend drastically on such changes, see [9].

Note, that the GNS states can be built only in the positive setting.

3 Clifford geometric approach

3.1 Nonsymmetric bilinear forms in Clifford algebras

To be able to describe the same structure as in the previous section, it is necessary to use a precision of the term Clifford algebra which might be called Clifford algebra of multivectors according to [38] or quantum Clifford algebras [20]. A detailed treatment can be found in [17, 19, 20], we give only the notations and results necessary for our models. Since a Clifford algebra is—in a functorial sense—the natural algebra of a quadratic space (V, Q) , a pair of a linear space V and a quadratic form Q . One can define a functor CL from the category of quadratic spaces into the category of unital associative algebras alg. But bilinear forms associated with quadratic forms are necessarily symmetric if the characteristic of the base field is not equal to 2. The polar bilinear form of a quadratic form Q is defined with $x, y \in V$ as

$$2B_p(x, y) := Q(x + y) - Q(x) - Q(y). \quad (16)$$

The Clifford map $\gamma : V \hookrightarrow CL$ results then in

$$\gamma_x \gamma_y + \gamma_y \gamma_x \simeq xy + yx = 2B_p(x, y), \quad (17)$$

where we identify $x \in V$ and $\gamma_x \in CL$. The Clifford product of elements from V , written without sign, can be decomposed as [39]

$$\gamma_x \gamma_y = B_p(x, y) + \gamma_x \wedge \gamma_y, \quad (18)$$

where the wedge is an antisymmetric Grassmann product. In writing this decomposition, we have used an isomorphism of the linear spaces FCL and $F \bigwedge V$, the linear space of Grassmann multiforms. This can be used to construct the Clifford algebra directly from the Grassmann algebra via Chevalley deformation [8]. The so defined product on V can be lifted on the whole algebra CL , see (20).

Moreover, we are no longer restricted to a symmetric bilinear form. One can define the left contraction on V

$$\gamma_x \lrcorner \gamma_y = B(x, y) \simeq \langle x | y \rangle \quad (19)$$

in a completely arbitrary way. $x \lrcorner$ is then $\in V^*$ and $\gamma_x \lrcorner$ its image in CL , see [37, 14, 12, 15, 1, 13, 16, 33]. Contrary to the associative wedge product, the contraction or

inner product is nonassociative. For multiforms we have to define with $x_i, y_j \in V$ and dropping from now on the injection $\gamma : V \hookrightarrow CL$

$$\begin{aligned}
i) \quad & x_i \lrcorner x_j = B(x_i, x_j) \simeq \langle x_i \mid x_j \rangle \\
ii) \quad & x_i \lrcorner (y_0 \wedge \dots \wedge y_n) = \sum_{r=0}^n (-1)^r (x_i \lrcorner y_r) (y_0 \wedge \dots \wedge y_{r-1} \wedge y_{r+1} \wedge \dots \wedge y_n) \\
iii) \quad & (x_1 \wedge \dots \wedge x_n) \lrcorner (y_1 \wedge \dots \wedge y_m) = x_1 \lrcorner (x_2 \lrcorner (\dots (y_1 \wedge \dots \wedge y_m) \dots)) \quad n \leq m.
\end{aligned} \tag{20}$$

We have however to remark, that the grading inherited from the Grassmann multivector structure is not unique. One could also define another contraction, which is symmetric and put the antisymmetric part into the Grassmann bi-form

$$\begin{aligned}
xy &= x \lrcorner_B y + x \wedge y = B(x, y) + x \wedge y \\
&= x \lrcorner_G y + x \dot{\wedge} y = G(x, y) + x \dot{\wedge} y.
\end{aligned} \tag{21}$$

Here $B = G + F$ is decomposed into symmetric and antisymmetric part, we have thus

$$x \dot{\wedge} y = F(x, y) + x \wedge y, \tag{22}$$

which clearly exhibits the different gradings. It was shown, that exactly this change of multivector structure is done when normal ordering, that is Wick ordering w.r.t. F , is performed in quantum (field) theory [24]. Even in nonlinear, resp. self-interacting, theories no singular additional terms arise when this change is treated algebraically correct [12].

In the sequel, we will give the bilinear forms explicitly in terms of a basis and use either the common wedge multiforms or monomials in Clifford generators. The Clifford monomials constitute a further possibility to introduce a grading on CL .

3.2 Fermi oscillator in Clifford terms

We start from a two dimensional ($n = 2$) hyperbolic space H , where we introduce the following particular basis $\{e_i\} = \{e_1, e_2\}$ and symmetric(!) bilinear form

$$[B(e_i, e_j)] = \left[\frac{1}{2} \delta_{i, n+1-i} \right] = \begin{vmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{vmatrix}. \tag{23}$$

We construct furthermore the Clifford algebra $CL(H, B)$, which leads to the commutation relations identifying from now on $\gamma_{e_i} \simeq e_i$ and $\mathbb{1} \simeq 1$

$$\begin{aligned}
e_i e_j + e_j e_i &= B(e_i, e_j) + B(e_j, e_i) + e_i \wedge e_j + e_j \wedge e_i \\
&= 2B(e_i, e_j).
\end{aligned} \tag{24}$$

If we identify e_1 with a and e_2 with $a^\dagger$, we notice that this algebra is isomorphic to the CAR algebra of the previous section. The relations then read

$$\begin{aligned}
aa^\dagger + a^\dagger a &= B(e_1, e_2) + B(e_2, e_1) = \frac{1}{2} + \frac{1}{2} = 1 \\
aa &= B(e_1, e_1) = 0 = B(e_2, e_2) = a^\dagger a^\dagger,
\end{aligned} \tag{25}$$

where the second line holds because of $a \wedge a = 0 = a^\dagger \wedge a^\dagger$. To establish the full correspondence, we need to define the $\dagger$ antiautomorphism. This can be done using the main antiinvolution $\sim$ of the Clifford algebra, also called reversion, defining the transformation $\underline{d}$ with help of the

$$p_i := (e_i - e_{n+1-i}) \quad (26)$$

as

$$\underline{d}(X) := \prod_{i=1}^{m=n/2=\dim V/2} (-p_i) X \tilde{p}_i = (-)^m p_1 p_2 \dots p_m X \tilde{p}_m \dots p_1. \quad (27)$$

It follows $\underline{d}(XY) = \underline{d}(Y)\underline{d}(X)$. In our case, we have

$$\begin{aligned} \underline{d}(e_1) &= e_2 & \Leftrightarrow & \underline{d}(a) = a^\dagger \\ \underline{d}(e_2) &= e_1 & \underline{d}(a^\dagger) &= a. \end{aligned} \quad (28)$$

Contrary to the ECAR case, where we had a correspondence of $FCAR$ and the dual elements $FCAR^*$, where each of the later constitutes a linear form, we have a canonical choice for a linear form in the Clifford algebra. We may define the scalar part $\langle \cdot \rangle_0$ of a Clifford number by projecting onto the field of scalars embedded in CL :

$$\begin{aligned} \langle \cdot \rangle_0 &: CL \mapsto \mathbb{K} \\ \langle A \rangle_0 &= \langle A \rangle_0^\wedge = \langle \alpha \mathbb{1} + \alpha_{ij} e_i \wedge e_j + \dots \rangle_0 = \alpha. \end{aligned} \quad (29)$$

We drop the wedge superscript when no confusion can occur which wedge is used to expand the Clifford numbers. Hence, the scalar part picks out the coefficient in front of the identity when A is expanded into *Clifford multivectors* w.r.t. *a certain Grassmann product*.

Now, the states ω_ν of the ECAR algebra shall be related to the single linear form of the Clifford algebra. We therefore use the representation theory of algebras. An irreducible, faithful representation can be obtained by a left regular representation of the simple algebra on a minimal left ideal generated by a primitive idempotent element. A good choice is either $P_{F^*} = aa^\dagger = e_1 e_2$, or $P_F = a^\dagger a = e_2 e_1$ because of

$$\begin{aligned} aa^\dagger aa^\dagger &= -a^2 (a^\dagger)^2 + aa^\dagger = aa^\dagger \\ a^\dagger aa^\dagger a &= -(a^\dagger)^2 a^2 + a^\dagger a = a^\dagger a. \end{aligned} \quad (30)$$

Furthermore, the ECAR relations show that the two projectors are a decomposition of unity

$$\mathbb{1} = P_F + P_{F^*} = aa^\dagger + a^\dagger a. \quad (31)$$

We can define now the Fock and dual Fock vacua by letting

$$\begin{aligned} \omega_F(X) &:= \langle 2P_F X \rangle_0 \\ \omega_{F^*}(X) &:= \langle 2P_{F^*} X \rangle_0, \end{aligned} \quad (32)$$

which are identical with the above defined forms, as can be seen by calculating their values on the generators. Since the factor 2 stems from the ungeometrical normalization of the ECAR relation, we would prefer the geometrical appropriate relation

$$\hat{a}_i \hat{a}_j^\dagger + \hat{a}_j^\dagger \hat{a}_i = 2\delta_{ij} \mathbb{1} = 2B_p(\hat{e}_i, \hat{e}_{n+1-j}), \quad (33)$$

remembering that $a_i^\dagger = e_{n+1-i}$. The ν -dependent vacua are then given as in (12) by

$$\begin{aligned} \omega_\nu(X) &:= v\omega_F(X) + (1-\nu)\omega_{F^*}(X) \\ &= \langle [\nu 2P_F + (1-\nu)2P_{F^*}]X \rangle_0. \end{aligned} \quad (34)$$

The grading operator this time, formulated within the algebra, is necessarily ν -dependent, because of the requirement $\omega_\nu(Q) = 0$ and found to be

$$Q := \left(\nu - \frac{1}{2}\right) - e_1 \wedge e_2 = \nu \mathbb{1} - aa^\dagger. \quad (35)$$

All results of the ECAR theory can be obtained now in purely Clifford algebraic terms, e.g. the μ -matrices as $[\mu_{ij}^\nu(A)] \simeq [\langle [\nu P_F + (1-\nu)P_{F^*}]d(E_i)AE_j \rangle_0]$ where the monomials E_i run in $\{\mathbb{1}, e_1e_2, e_1, e_2\}$. However, since we do not need a representation, but used the basis only to show the equivalence to the above discussed C^* -algebraic description, we need not bother about further transformations and positivity requirements.

Since we have up to now hardly done more than to reformulate the ECAR results, we have to face the question why we prefer a Clifford geometric approach. Even the correspondence to a single linear form, sic a trace, is common to physicists, who are used to working with matrix representations. Every state on a finite dimensional real or complex (E)CAR algebra can be written as

$$\omega_\rho(X) = \frac{1}{\dim V} \text{Tr}(\rho\pi(X)), \quad (36)$$

where V is the representation space, $\pi : (E)CAR \mapsto \text{End}(V)$ a representation and ρ the density matrix. However, a representation is implicitly needed in such a construction. In the next subsection we utilize the Clifford algebra in a new and geometric way. This step, taking the Clifford geometric character fully into account, is the doorway to generalize the situation thereafter *beyond* ECAR possibilities.

3.3 Clifford geometric Fermi oscillator

In the previous subsection we have modelled the same situation as in the ECAR algebra by simply translating them into Clifford terms. It is however clear, that the full power of a mathematical tool can only be achieved if its generic abilities are used. Hence, we will give a further approach to the Fermi oscillator which relies fully on Clifford geometric methods.

Let B_ν be a bilinear form on the space V , ($\dim V = 2$) which generates $CL(V, B_\nu)$. In a distinguished basis $\{e_i\} = \{e_1, e_2\}$ we have

$$\begin{aligned} [B_\nu] &= \begin{vmatrix} 0 & \nu \\ 1-\nu & 0 \end{vmatrix} = \begin{vmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{vmatrix} + \begin{vmatrix} 0 & -\frac{1}{2} + \nu \\ \frac{1}{2} - \nu & 0 \end{vmatrix} \\ &= [G] + [F_{-1/2+\nu}] = [G] + [F_\nu]. \end{aligned} \quad (37)$$

The normalization is chosen in such a way, that ν will have the same values as above for positive solutions, $\nu \in]0, 1[$. Defining a contraction $\lrcorner_\nu$ on V and lifting it to multivector arguments as in (20), we come up with the Clifford algebra $CL(V, B_\nu)$. Contraction, Clifford and wedge products are related by

$$\begin{aligned} x \lrcorner_\nu y &\equiv x \lrcorner_{B_\nu} y = B_\nu(x, y) \\ xy &= x \lrcorner_\nu y + x \wedge y \\ &= x \lrcorner_G y + x \lrcorner_{F_{-1/2+\nu}} y + x \wedge y \\ &= x \lrcorner_G y + x \hat{\wedge} y. \end{aligned} \tag{38}$$

Obviously, $\wedge$ and $\hat{\wedge}$ define different multivector structures in CL . Indeed, we had already mentioned that CL is only $\mathbb{Z}_2$ -graded and bears *no natural* multivector structure. In other words, CL does not depend on the $\mathbb{Z}_n$ -grading and we have a Clifford algebra isomorphism, that is the Wick theorem [12, 13, 20, 24], $CL(V, B_\nu) \simeq CL(V, G)$, where G is symmetric and does correspond directly to a quadratic form or say a “quantization”. This can be seen by inspection of the commutator relations which turn out to be independent of ν ($n = \dim V$)

$$\begin{aligned} e_i e_j + e_j e_i &= B_\nu(e_i, e_j) + B_\nu(e_j, e_i) \\ &= 2G(e_i, e_j) = \delta_{i, n+1-j} \\ &= Q(e_i + e_j) - Q(e_i) - Q(e_j). \end{aligned} \tag{39}$$

This yields immediately the identification $a = e_1, a^\dagger = e_2$. If we would like to insist on symmetry, we should now shift ν to $\nu' = -1/2 + \nu$. But to be able to compare the results with previous sections, we remain with ν .

Since we noted, that the scalar-part projection depends on the multivector structure, which is now parameterized by ν , we have an intrinsic way opened to describe vacuum states. However, this is now not a question of a dual-isomorphism connecting V and V^* , positivity and a statistical interpretation, but simply a matter of the geometry emerging from an additional antisymmetric part in the contraction. Moreover, we showed in [15, 13] that F equals the propagator of the theory. We have thus found a constructive way to relate the dynamics i.e. propagator data to representations.

To demonstrate the equivalence of $\langle \dots \rangle_0^\wedge$ –scalar-part of $CL(V, B_\nu)$ without any Fock-projectors involved– and ω_ν we calculate the values of *the single state* $S^0(CL(V, B_\nu))$ on the generators (the linear form taken element-wise)

$$\langle \{ \mathbb{1}, aa^\dagger, a, a^\dagger \} \rangle_0^\wedge = \{ 1, \nu, 0, 0 \}, \tag{40}$$

compare with eqn. (5). Since the Clifford product and the anticommutation relations are ν -independent, we have established an algebra isomorphism. Since we had implemented the $\dagger$ via a Clifford reversion followed by a linear transformation $\underline{d}$, we have in fact constructed a $*$ -isomorphism. One should remark however that the Clifford reversion does *not* respect in general the multivector structure. This can be seen from

$$\begin{aligned} (e_1 \wedge e_2)^\sim &= [e_1 e_2 - B_\nu(e_1, e_2)]^\sim \\ &= e_2 e_1 - B_\nu(e_1, e_2) = e_2 \wedge e_1 + B_\nu(e_2, e_1) - B_\nu(e_1, e_2) \\ &= -e_1 \wedge e_2 - 2F_{1/2-\nu}(e_1, e_2). \end{aligned} \tag{41}$$

On the other hand, the dotted wedge $\hat{\wedge}$ defined in (38) is stable under reversion [16]. This motivated the normal ordering procedure in QM and QFT [12, 13]. However, one has *carefully to distinguish* then the scalarpart projections $\langle \dots \rangle_0^{\hat{\wedge}}$ and $\langle \dots \rangle_0^{\wedge}$, which result in entirely different states. The later constitutes the maximally mixed state w.r.t. $\wedge$

$$\langle \{\mathbb{1}, aa^\dagger, a, a^\dagger\} \rangle_0^{\hat{\wedge}} = \{1, \frac{1}{2}, 0, 0\} \Rightarrow \nu = \frac{1}{2}, \quad (\nu' = 0), \quad (42)$$

which of course also equals the usual *normal ordered* Fock state w.r.t. $\hat{\wedge}$. The state is positive if $\nu \in]0, 1[$, which is equivalently written as

$$-\det B_\nu > 0. \quad (43)$$

The relation between geometry and algebraic norms of such types were examined in [14]. In the cases $\nu = 0$ and $\nu = 1$, the base space has only a degenerated form, which does not allow the construction of a universal Clifford algebra. Finally, we could furthermore calculate once more, to show the equivalence to our previous results, the matrices $[\mu_{ij}^\nu(X)] = [\langle \underline{d}(E_i) X E_j \rangle_0^{\hat{\wedge}}]$. One should note, that within the Clifford algebraic approach, a representation does not become necessary, since all calculations can be performed within *abstract* algebra. Hence, we are not troubled with the positivity requirement to obtain representations in Hilbert spaces via the GNS construction.

3.4 Two degrees of freedom

As we have already said, the one dimensional system served as a learning field. In moving to the next dimension, we will however see some probably unexpected details. This system is furthermore able to describe physical situations, especially the occurrence of spin-zero and spin-one can be used for the purpose of bosonization.

The two-dimensional system also exhibits differences between the ECAR and Clifford approach, which goes beyond the simple restriction of some parameters to gain positivity. The coherence that *all* entities in the theory are formulated within *one* mathematical system, in the multivector Clifford geometric algebra, unfolds dependencies and relations which were in principle not visible in the former ECAR setting. This is a further example of the usability of geometric quantum Clifford algebras as claimed by Hestenes [29].

3.4.1 ECAR results

Since we fear having already overdone the obvious in the treatment of the one-dimensional system, we will give in this section only the definitions and results. Most of them can be found in [31, 32].

We define the $U(2)$ -ECAR algebra generated by $\{a_\alpha, a_\beta^\dagger, \mathbb{1}, S_k, Q\}$, $\alpha, \beta \in \{1, 2\}$, $k \in \{1, 2, 3\}$ via the relations

$$\begin{aligned} \{a_\alpha, a_\beta\} &= 0 = \{a_\alpha^\dagger, a_\beta^\dagger\} \\ \{a_\alpha, a_\beta^\dagger\} &= \delta_{\alpha, \beta} \mathbb{1}. \end{aligned} \quad (44)$$

Table 1.				
No:	$A \in ECAR$	s_3	$s(s+1)$	q
g_1	$\mathbb{1}$	0	0	0
g_2	$\frac{1}{2}(a_1 a_1^\dagger + a_2 a_2^\dagger)$	0	0	0
g_3	$a_1 a_2 a_2^\dagger a_1^\dagger$	0	0	0
g_4	$a_1 a_2^\dagger$	1	2	0
g_5	$\frac{1}{2}(a_1 a_1^\dagger - a_2 a_2^\dagger)$	0	2	0
g_6	$a_2 a_1^\dagger$	-1	2	0
g_7	a_1	$\frac{1}{2}$	$\frac{3}{4}$	1
g_8	$a_1 a_2 a_2^\dagger$	$\frac{1}{2}$	$\frac{3}{4}$	1
g_9	a_2	$-\frac{1}{2}$	$\frac{3}{4}$	1
g_{10}	$a_2 a_1 a_1^\dagger$	$-\frac{1}{2}$	$\frac{3}{4}$	1
g_{11}	$a_2^\dagger$	$\frac{1}{2}$	$\frac{3}{4}$	-1
g_{12}	$a_1 a_1^\dagger a_2^\dagger$	$\frac{1}{2}$	$\frac{3}{4}$	-1
g_{13}	$a_1^\dagger$	$-\frac{1}{2}$	$\frac{3}{4}$	-1
g_{14}	$a_2 a_2^\dagger a_1^\dagger$	$-\frac{1}{2}$	$\frac{3}{4}$	-1
g_{15}	$a_1 a_2$	0	0	2
g_{16}	$a_1^\dagger a_2^\dagger$	0	0	-2

Table 1.: Eigenvectors used as basis states and their $U(2)$ quantum numbers

Furthermore, we demand an action of the group $U(2)$, if the complex ground field is used, by the requirements

$$\begin{aligned}
[Q, S_k] &= 0, \quad [S_k, S_l] = i\epsilon_{klm} S_m \\
S^\dagger &= S, \quad Q^\dagger = Q \\
[S_k, a_\alpha] &= \sigma_k^{\alpha\beta} a_\beta, \quad [S_k, a_\alpha^\dagger] = \hat{\sigma}_k^{\alpha\beta} a_\beta^\dagger \\
[Q, a_\alpha] &= +a_\alpha, \quad [Q, a_\alpha^\dagger] = -a_\alpha^\dagger,
\end{aligned} \tag{45}$$

where $\sigma_k^{\alpha\beta}$ are the Pauli-matrices and $\hat{\sigma}_k = -\bar{\sigma}_k = (-)^k \sigma_k$ (the relations are not independent). Since we do not fix any dependency on the S_k, Q as functions of the $\{a_\alpha, a_\alpha^\dagger, \mathbb{1}\}$, we have defined an Extended CAR algebra, [11].

A maximal set of pair-wise commuting group elements is given by Q, S_3 and $\vec{S}^2 = \sum_{k=1}^3 S_k^2$. We feel very uncomfortable with this notation, because S_k will be a multivector aggregate within the Clifford algebra and not a ‘vector’, but we use it because of readability and since it is a standard.

Additionally, we define a distinguished basis $\{g_i\}$ in terms of polynomials in the Clifford generators, which are chosen to be eigenvectors of the commuting operators $S_3, \vec{S}^2, Q$, with eigenvalues $s_3, s(s+1), q$. This is summarized in table 1, see [31].

The linear space spanned by $\{g_1, g_2, g_3\}$ contains the candidates for vacuum states. To be precise, $\{g_1, g_2, g_3\}$ are in the kernel of the map $\omega : FA \mapsto \mathbb{K}$. One may note the occurrence of a spin-one triplet ($s(s+1) = 2$) which of course has zero

expectation values in Fock-space, of two spin one-half “quartets” ($s(s+1) = 3/4$) of opposite charge or “particle number” and two spin zero eigenvectors of “particle number” $q = \pm 2$.

In [31, 32] the vacuum states were deduced from special conditions (given below, (46)) on this basis. However, one might notice that these states are defined only up to an additive, possibly complex constant e.g. $g_5 \simeq g'_5 = g_5 + c\mathbb{1}$, since the constant does not contribute to the commutator e.g. $[Q, g_5] \simeq [Q, g'_5] = [Q, g_5 + c\mathbb{1}]$. This fact spoils the claim in [31, 32] that one can conclude from $\langle \Omega | g_5 \Omega \rangle = \langle \Omega | \frac{1}{2}(a_1 a_1^\dagger - a_2 a_2^\dagger) \Omega \rangle = 0$ the validity of $\langle \Omega | a_1 a_1^\dagger \Omega \rangle = - \langle \Omega | a_2 a_2^\dagger \Omega \rangle (= \nu)$. We will nevertheless stay with this restriction to be able to compare ECAR and Clifford geometric results and impose the *normalizations* of the g_i given in table 1 as further constraints. However, we will see that the pair (ν, w) is sufficient to parameterize all vacuum states, but the ECAR and thereby the C^* -algebraic approach is then no longer able to relate them to other data, as to the propagator. This is a major drawback not apparent in the multivector Clifford algebraic framework.

With the basis of table 1 (including normalization), we conclude that vacuum states can be parameterized by two real variables ν, w since we have

$$\begin{aligned} \omega_{\nu w}(g_1) &= \omega_{\nu w}(\mathbb{1}) = 1 && \text{normalization of } \omega_{\nu w} \\ \omega_{\nu w}(g_5) &\Rightarrow \omega_{\nu w}(a_1 a_1^\dagger) = \omega_{\nu w}(a_2 a_2^\dagger) = \nu && \text{see remark above} \\ \omega_{\nu w}(g_3) &= \omega_{\nu w}(a_1 a_2 a_2^\dagger a_1^\dagger) = w \\ \omega_{\nu w}(g_4) &= \dots = \omega_{\nu w}(g_{16}) = 0 && \text{bec. of nontrivial eigenvalues. (46)} \end{aligned}$$

If we further require that $\omega_{\nu w}$ is a positive state, we have to restrict the parameters to

$$\begin{aligned} 1 &> \nu > w > 0 \\ w &> 2\nu - 1, \end{aligned} \tag{47}$$

which describes the interior of a simplex, see Fig. 1. It was demonstrated in [31, 32] that $\omega_{\nu w}$ can be decomposed into three extremal states, once more the Fock and dual Fock states with $\omega_F = \omega_{11}$, $\omega_{F^*} = \omega_{00}$ and a further uncommon state $\omega_E = \omega_{1/2 0}$. This yields the decomposition

$$\omega_{\nu w} = w\omega_F + (1 - 2\nu + w)\omega_{F^*} + 2(\nu - w)\omega_E. \tag{48}$$

Quasi free states, with vanishing higher correlations [7] turn out to be described by $\omega_{\nu \nu^2}$, a parabola in the ν - w -plane. It is interesting that Bogoliubov transformations which mix Fock and dual Fock states do therefore create correlations, but with no contribution of ω_E , see also discussion below.

We end now the discussion, further information on the dynamics and the thermodynamical behaviour generated by this type of system can be found in [31, 32]. However, one should keep in mind that the ECAR treatment is not able to relate the parameters ν, w to dynamical data.

3.4.2 Clifford geometric results

If we use any possible freedom in defining a contraction, the Clifford algebra provides a richer structure than the ECAR algebra. Since we want to stay with a correspondence $\{a_1, a_2, a_2^\dagger, a_1^\dagger\} = \{e_1, e_2, e_3, e_4\}$, we require that the symmetric part of the contraction of the $\{a_i\}$ and $\{a_j^\dagger\}$, leads to

$$\begin{aligned} \{a_\alpha, a_\beta\} &= 0 = \{a_\alpha^\dagger, a_\beta^\dagger\} \\ \{a_\alpha, a_\beta^\dagger\} &= \delta_{\alpha\beta} \mathbb{1}. \end{aligned} \quad (49)$$

Canonical quantization turns out to be nothing more than the specification of an appropriate quadratic form in use [15, 13]. The antisymmetric part of the contraction is *a priori* arbitrary; we set therefore

$$\begin{aligned} [a_\alpha \lrcorner_B a_\beta^\dagger] &= [e_i \lrcorner_B e_{j+2}] = \begin{vmatrix} q & r \\ s & t \end{vmatrix} \\ [a_\alpha^\dagger \lrcorner_B a_\beta] &= [e_{i+2} \lrcorner_B e_j] = \begin{vmatrix} -q & 1-s \\ 1-r & -t \end{vmatrix}, \end{aligned} \quad (50)$$

and note that from $\{a_\alpha, a_\beta\} = 0 = \{a_\alpha^\dagger, a_\beta^\dagger\}$ no additional information is obtained from the antisymmetric part, which finally yields in terms of Clifford generators

$$[B(e_i, e_j)] = \begin{vmatrix} 0 & u & q & r \\ -u & 0 & s & t \\ -q & 1-s & 0 & m \\ 1-r & -t & -m & 0 \end{vmatrix}. \quad (51)$$

A canonical grading operator may be defined by the identity operator on the space spanned by the $\{a_\alpha\}$ alone

$$Q' := a_1 a_1^\dagger + a_2 a_2^\dagger. \quad (52)$$

The vacuum state is defined as the scalarpart $\langle \dots \rangle_0^\wedge$. We have thus to renormalize Q' by a scalar additive constant. From

$$\langle Q \rangle_0^\wedge := \langle Q' + \alpha \mathbb{1} \rangle_0^\wedge = 0 \quad (53)$$

we deduce

$$\begin{aligned} -\alpha &= \langle Q' \rangle_0^\wedge = \langle e_1 e_4 + e_2 e_3 \rangle_0^\wedge \\ &= \langle B(e_1, e_4) + B(e_2, e_3) + e_1 \wedge e_4 + e_2 \wedge e_4 \rangle_0^\wedge \\ &= B(e_1, e_4) + B(e_2, e_3) = r + s, \end{aligned} \quad (54)$$

or

$$Q = a_1 a_1^\dagger + a_2 a_2^\dagger - (r + s) \mathbb{1} = a_1 \wedge a_1^\dagger + a_2 \wedge a_2^\dagger. \quad (55)$$

Observe, that the first expression for Q is independent of the specific used $\wedge$, but depends on a specific normalization $-(r+s)$, while the second is independent of the normalization, but depends on the specific used $\wedge$. From table 1, we know that $\{g_1, g_2, g_3\}$ might span the vacuum sector. A vacuum state ω_X should thereby fulfil $\omega_X(g_4) = \dots \omega_X(g_{16}) = 0$. We calculate $\langle \dots \rangle_0^\wedge$ on all basis elements $\{g_i\}$ to obtain

$$\begin{aligned} \{\langle g_i \rangle_0^\wedge\} &= \{1, \frac{1}{2}(r+s), um - tq + rs, q, \frac{1}{2}(r-s), t, \\ &\quad 0, 0, 0, 0, 0, 0, u, -m\}. \end{aligned} \quad (56)$$

Because of the choice of normalization in table 1, and the requirement that $\langle \dots \rangle_0^\wedge$ is a vacuum state, we have to set $\langle g_4 \rangle_0^\wedge = \langle g_5 \rangle_0^\wedge = \langle g_6 \rangle_0^\wedge = \langle g_{15} \rangle_0^\wedge = \langle g_{16} \rangle_0^\wedge = 0$, which yields

$$q = 0, \quad t = 0, \quad r = s, \quad u = 0, \quad m = 0. \quad (57)$$

The third relation is equivalent to the condition $\omega_{\nu w}(a_1 a_1^\dagger) = \omega_{\nu w}(a_2 a_2^\dagger) = \nu$ in (46). But the Clifford treatment uncovers then the relation of $\nu = r = s$ and $w = um - tq + rs = \langle g_3 \rangle_0^\wedge$. Since we have to fulfil (57) consistency is obtained only if we have $w = rs = \nu^2$ which leads to

$$\omega_{\nu \nu^2} = \langle \dots \rangle_0^\wedge, \quad q = t = u = m = 0, \quad r = s = \nu. \quad (58)$$

These are quasi free states and thereby almost trivial. The space of states is parameterized by one single real number ν , not a pair (ν, w) . To provide a set of positive states, ν has to be restricted to $]0, 1[$.

The Clifford algebraic examination of the ECAR results unveils therefore an inconsistency, which breaks up the relation of contraction data and the parameterization of vacuum states and quantization. Below, we discuss the connection of the propagator, contraction and these parameters in Clifford geometric terms, which is thus *beyond* the ability of ECAR methods.

Since we want to look for more general situations, we use the freedom to renormalize the basis $\{g_i\}$, which is defined only up to constants $g'_i = g_i + c_i \mathbb{1}$. If we set

$$\begin{aligned} g'_4 &= g_4 - q \\ g'_5 &= g_5 - \frac{1}{2}(r-s) \\ g'_6 &= g_6 - t, \\ g'_{15} &= g_{15} - u, \\ g'_{16} &= g_{16} + m, \end{aligned} \quad (59)$$

we cancel the unintentional expectation values. We drop the prime thereafter. The vacuum states can then be parameterized by two real numbers ν, w , which are of course functions of the parameters $q, r, s, t; u, m$. One obtains from g_2 and g_3

$$\begin{aligned} \langle g_2 \rangle_0^\wedge &= \nu = \nu(q, r, s, t; u, m) = \frac{1}{2}(r+s) \\ \langle g_3 \rangle_0^\wedge &= w = w(q, r, s, t; u, m) = um - tq + rs. \end{aligned} \quad (60)$$

If we set $r = -s + 2\nu$, we remain with $w = um - tq - s^2 + 2s\nu$, which does posses the solutions

$$s = \nu \pm \sqrt{\nu^2 - tq + um - w} \quad \# \text{ variables} = 4, \quad (q, t, u, m). \quad (61)$$

To be able to relate this results with more conventional approaches, we remark that the propagator, also denoted by F , of the theory was shown to be equivalent to the antisymmetric part of B in [15, 12, 13]. This can be seen as follows

$$\begin{aligned} F_{ij} &= \langle \frac{1}{2}[a_i, a_j^\dagger] \rangle_0^\wedge = \frac{1}{2} \langle a_i a_j^\dagger - a_j a_i^\dagger \rangle_0^\wedge \\ &= \frac{1}{2} \langle B(e_i, e_{n+1-j}) - B(e_{n+1-j}, e_i) + e_i \wedge e_{n+1-j} - e_{n+1-j} \wedge e_i \rangle_0^\wedge \\ &= \langle F(e_i, e_{n+1-j}) + e_i \wedge e_{n+1-j} \rangle_0^\wedge \\ &= F(e_i, e_{n+1-j}), \end{aligned} \quad (62)$$

and hence

$$[F_{ij}] = \langle \frac{1}{2}[a_i, a_j^\dagger] \rangle_0^\wedge = \begin{vmatrix} -r + 1/2 & q \\ t & -s + 1/2 \end{vmatrix} \quad (63)$$

The solutions obtained in (61) yield thus a direct and *constructive* relation between the matrix elements of the propagator $(q, r, s, t; u, m)$ and the parameterization of the corresponding vacua (ν, w) .

Since we have to compare our results with ECAR algebraic ones over the complex number field, we have to take the $(q, r, s, t; u, m)$ parameters as complex numbers. From

$$[F_{\alpha\beta}]^\dagger = \langle \frac{1}{2}([a_\alpha, a_\beta^\dagger])^\dagger \rangle_0^\wedge = \langle \frac{1}{2}[a_\beta, a_\alpha^\dagger] \rangle_0^\wedge = [F_{\beta\alpha}] \quad (64)$$

or in matrix form

$$\begin{vmatrix} -\bar{r} + 1/2 & \bar{t} \\ \bar{q} & -\bar{s} + 1/2 \end{vmatrix} = \begin{vmatrix} -r + 1/2 & t \\ q & -s + 1/2 \end{vmatrix} \quad (65)$$

we conclude, that (q, r, s, t) have to be real numbers. From

$$\begin{aligned} u^\dagger &= \langle g_{15}^\dagger \rangle_0^\wedge = \langle (a_1 a_2)^\dagger \rangle_0^\wedge = - \langle a_1^\dagger a_2^\dagger \rangle_0^\wedge \\ &= - \langle g_{16} \rangle_0^\wedge = m \end{aligned} \quad (66)$$

we obtain that $\Delta_0 := um = uu^\dagger \geq 0$ and analogously from $q^\dagger \simeq g_4^\dagger = g_6 \simeq t$, $q = q^\dagger = t = t^\dagger$ the relation $\Delta_1 := qt = qq^\dagger = q^2 \geq 0$. The requirement of hermiticity has thus furthermore restricted the parameters. The Δ_i are shifts induced by the normalization of spin-zero and spin-one eigenvectors. From (61) we remain with the ‘gap-equation’ (the name will become clear later)

$$s = \nu \pm \sqrt{\nu^2 - \Delta_1 + \Delta_0 - w}. \quad (67)$$

The propagator can finally be written as

$$F_{\alpha\beta}(\nu, w, \Delta_1 - \Delta_0, q) = \begin{vmatrix} -\nu \pm \sqrt{\nu^2 - \Delta_1 + \Delta_0 - w} & q \\ q & -\nu \mp \sqrt{\nu^2 - \Delta_1 + \Delta_0 - w} \end{vmatrix} \quad (68)$$

Let us look more closely at the vacuum sector obtained in our $U(2)$ model. We had already seen, that the space of all positive vacua is an affine simplex –thereby convex–, spanned by three extremal states: the Fock state $\omega_F = \omega_{11}$, the dual Fock state $\omega_{F^*} = \omega_{00}$ and an state $\omega_E = \omega_{1/20}$. A general state was decomposed as

$$\omega_{\nu w} = w\omega_F + (1 - 2\nu + w)\omega_{F^*} + 2(\nu - w)\omega_E. \quad (69)$$

Since Bogoliubov transformations do mix Fock and dual Fock states, they generate an edge of the simplex, which might also be parameterized as

$$\omega_{BV} = \omega_{1-\rho 1-\rho} = \rho\omega_{F^*} + (1 - \rho)\omega_F, \quad \rho \in]0, 1[. \quad (70)$$

The full space of positive vacua is then a convex combination of ω_{BV} and ω_E with $\lambda \in]0, 1[$

$$\begin{aligned} \omega_{\nu w} = \omega'_{\lambda\rho} &= \lambda\omega_{BV} + (1 - \lambda)\omega_E \\ &= \lambda(1 - \rho)\omega_F + \lambda\rho\omega_{F^*} + (1 - \lambda)\omega_E. \end{aligned} \quad (71)$$

If we now fix a ρ and look at the line of vacua parameterized by λ , we get a classical bifurcation diagram for the solutions of (61) most easily seen if the Δ_i are zero, see Fig 1. The quasi free states given by $\nu^2 = w$ do constitute the borderline of different phases. Since quasi free states are defined to have no higher correlations, they separate areas which do posses higher correlations of possibly different signs. In our case, one area has an attractive force and an ordered phase, while the other one has repulsive character. In the area between ω_{BV} and $\omega_{\nu \nu^2 - \Delta_1 + \Delta_2}$ one has a typical gap-equation, coming from the two real roots of eqn. (67), where between $\omega_{\nu \nu^2 - \Delta_1 + \Delta_2}$ and the edge state ω_E this type of solution can not occur in (67) since s has to be real.

The C^* -algebraic ECAR solution of the QFT hierarchy equations of BCS-theory can be found in [31, 32], which supports our terms: vacuum, phase transition, ordered phase, gap-equation, etc. Copper pairs should be identified with spin-0 states of particle number –grading– 2. See also the following discussion.

4 Discussion

4.1 Relation to Bogoliubov–Valatin transformations

We want to introduce here the common notation of a Fock vacuum more explicitly. This will show the equivalence of one of the above derived vacua to this particular state, but also the restricted validity of the Fock approach. Especially in QFT, where Haag’s theorem rules out this possibility.

Let $|0\rangle_F$ ($|0\rangle_{F^*}$) be a (dual) Fock vector state which is subjected to the conditions

$$\begin{aligned} \prod_{i \in I} a_i |0\rangle_F &= 0 \\ (\prod_{i \in I} a_i^\dagger |0\rangle_{F^*} &= 0) \end{aligned} \quad (72)$$

where I contains non-empty ordered subsets of cardinality less than or equal to N .

The question is, whether this definition is equivalent to our above description of Fock and dual Fock states. This can be seen as follows: We calculate, introducing the shorthand ${}_F \langle 0 | X | 0 \rangle_F = \langle X \rangle_F$, the expectation value

$$0 = \langle a_i^\dagger a_i \rangle_F = \langle \mathbb{1} - a_i a_i^\dagger \rangle_F = 1 - \begin{cases} r & i = 1 \\ s & i = 2 \end{cases}. \quad (73)$$

Now, $r = s = \nu$ according to (58,60) and we end up with

$$\nu = 1. \quad (74)$$

From

$$\begin{aligned} 0 &= \langle a_2^\dagger a_1^\dagger a_1 a_2 \rangle_F = \langle a_2^\dagger a_2 - \mathbb{1} + a_1^\dagger a_1 + a_1 a_2 a_2^\dagger a_1^\dagger \rangle_F \\ &= 0 - 1 + 0 + w, \end{aligned} \quad (75)$$

we get

$$w = 1. \quad (76)$$

We conclude therefore, that the Fock vacuum is given in our framework as

$$\omega_F = \omega_{\nu w} \Big|_{\nu=1 \ w=1} = \omega_{11}, \quad (77)$$

whereas the dual-Fock state can be deduced to be given along the same lines as

$$\omega_{F^*} = \omega_{\nu w} \Big|_{\nu=0 \ w=0} = \omega_{00}. \quad (78)$$

The state space of the (dual) Fock vacuum is spanned by the following vectors

$$\begin{aligned} \mathcal{V}_F &= \{ \mathbb{1} | 0 \rangle_F, a_1^\dagger | 0 \rangle_F, a_2^\dagger | 0 \rangle_F, a_1^\dagger a_2^\dagger | 0 \rangle_F \} \\ \mathcal{V}_{F^*} &= \{ \mathbb{1} | 0 \rangle_{F^*}, a_1 | 0 \rangle_{F^*}, a_2 | 0 \rangle_{F^*}, a_1 a_2 | 0 \rangle_{F^*} \}. \end{aligned} \quad (79)$$

Both spaces are four dimensional, but yield different matrix representations for the group generators e.g. N . This shows explicitly their inequivalence.

In the case of the new edge state, we turn around the direction of inference to deduce from

$$\omega_E = \omega_{\nu w} \Big|_{\nu=1/2 \ w=0} = \omega_{1/2 0} \quad (80)$$

the conditions

$$a_1 a_2 |0\rangle_E = 0 = a_2^\dagger a_1^\dagger |0\rangle_E. \quad (81)$$

The state space of this edge is generated by

$$\mathcal{V}_E = \{\mathbb{1}, a_1, a_2, a_1^\dagger, a_2^\dagger, \frac{1}{2}(a_1 a_1^\dagger - a_2 a_2^\dagger), a_1 a_2^\dagger, a_2 a_1^\dagger\} |0\rangle_E. \quad (82)$$

Note, that neither a_i nor $a_i^\dagger$ annihilate this vacuum state. Hence, a linear Bogoliubov-Valatin like transformation $c_i = f(a_i, a_j^\dagger) = \alpha_i a_i + \beta_j a_j^\dagger$ cannot reach this type of vacuum state. Bogoliubov-Valatin vacua remain on a single line in our vacuum plane given by $\omega_{BV} = \omega_{1-\rho} 1 - \rho$, see Fig 1. To relate our results to the literature, we investigate this connection a little more.

In the theory of superconductivity, Bogoliubov [6], Valatin [43] as Holstein and Primakoff [30] introduced new quasi particles $c, c^\dagger$ via the definition

$$\begin{aligned} c_1 &:= \sqrt{1-h} a_1 - \sqrt{h} a_2^\dagger \\ c_2 &:= \sqrt{1-h} a_2 + \sqrt{h} a_1^\dagger. \end{aligned} \quad (83)$$

The transformation is canonical since it fulfils

$$\{c_i, c_k^\dagger\} = \delta_{ik} \mathbb{1}, \quad \{c_i, c_k\} = 0 = \{c_i^\dagger, c_k^\dagger\}. \quad (84)$$

Furthermore, a new quasi-particle Fock vacuum $|0\rangle_{BV} = [\sqrt{1-h} + \sqrt{h} a_1^\dagger a_2^\dagger] |0\rangle_F$ is introduced, demanding

$$c_i |0\rangle_{BV} = 0 \quad \forall i. \quad (85)$$

Obviously, $|0\rangle_{BV}$ and $|0\rangle_F$ are quite different vector states of the underlying state space. Furthermore, none of these states are vacuum states w.r.t. the alternate choice of creation and annihilation operators. One finds

$$\begin{aligned} c_1 |0\rangle_F &= \sqrt{1-h} a_1 + \sqrt{h} a_2^\dagger |0\rangle_F = \sqrt{h} |0, 1\rangle_F \neq 0 \\ a_1 |0\rangle_{BV} &= a_1 [\sqrt{1-h} + \sqrt{h} a_1^\dagger a_2^\dagger] |0\rangle_F = \sqrt{h} |0, 1\rangle_F \neq 0 \end{aligned} \quad (86)$$

etc. in an obvious notation. From this consideration, it is quite clear, that the notation of a Fock vacuum, i.e. annihilator on a groundstate equals zero, is highly dependent on the chosen set of creation and annihilation operators. One cannot speak in this terms of a *vacuum* but only of a vacuum w.r.t. $a, a^\dagger$ or $c, c^\dagger$ etc. Moreover, the naive mixture of creation and annihilation operators spoils particle number conservation w.r.t. the old variables, as also the state $|0\rangle_{BV}$ has no definite a -particle number. In consequence, this means that electrons and Cooper pairs cannot be sharply measured simultaneously in such a framework. Moreover, even if we are able to mimic the Bogoliubov-Valatin results, strictly speaking this state is not a vacuum in our sense.

An algebraic treatment of states as given above does mix states and representations in favour of operators and is thus able to handle mixed states without the described

problems, see [31, 32]. However, much more important is the fact that Bogoliubov–Valatin transformations can only create correlations due to a mixture of Fock and dual Fock states and fail to be able to give the full possibilities of the three extremal, or say, pure cases. No λ mixture can appear as in (71). The algebraic setting based on GNS techniques in the positive case and extended to indefinite cases by Clifford algebraic methods is thus more general than Bogoliubov-Valatin transformations.

4.2 BCS propagator according to Fetter Walecka

To be able to compare our results to the ones usually given in literature, we derive some further identities. We extend our theory in the sense that we add a new mode index k to our fields, as we introduce an independent copy of the algebra for every mode. This is not harmful, but makes all of our real parameters functions of k . The BCS groundstate is given as [4]

$$|0\rangle_{BCS} = \Pi_k [\sqrt{1-h_k} + \sqrt{h_k} a_{k1}^\dagger a_{-k2}^\dagger] |0\rangle_F. \quad (87)$$

The connection to the Bogoliubov-Valatin transformation is then given by

$$\begin{aligned} c_{k1} &:= u_k a_{k1} - v_k a_{-k2}^\dagger \\ c_{k2} &:= u_k a_{k2} + v_k a_{-k1}^\dagger, \end{aligned} \quad (88)$$

compare (83), with the new vacuum defined as Fock vacuum of the quasi particles. Comparing (88) and (83) we get

$$\begin{aligned} u_k &= \sqrt{1-h_k} \\ v_k &= \sqrt{h_k}. \end{aligned} \quad (89)$$

Comparing these quantities with our data, i.e. the vacuum parameters, we obtain the u_k and v_k in terms of these parameters. From the Bogoliubov-Valatin vacuum $\omega_{1-\rho_k} 1-\rho_k$ we have

$$\begin{aligned} c_{k1} &= \sqrt{\frac{\rho_k-1}{2\rho_k-1}} e_{k1} - \sqrt{\frac{\rho_k}{2\rho_k-1}} e_{-k3} \\ c_{k2} &= \sqrt{\frac{\rho_k-1}{2\rho_k-1}} e_{k2} + \sqrt{\frac{\rho_k}{2\rho_k-1}} e_{-k4}, \end{aligned} \quad (90)$$

and thus

$$\begin{aligned} \nu_k = w_k &= 1 - \rho_k \\ u_k &= \sqrt{\frac{\rho_k-1}{2\rho_k-1}} \\ v_k &= \sqrt{\frac{\rho_k}{2\rho_k-1}}, \end{aligned} \quad (91)$$

iff we set the shifts $u = m = 0$. From $\nu_k = w_k = 1 - \rho_k$ we can derive all of our propagator data etc. in terms of ρ_k and finally in terms of the u_k, v_k parameters of the

Bogoliubov-Valatin transformation. We can thus identify the results given in literature with those given by Clifford algebraic considerations.

We follow Fetter and Walecka (FW) [25]. The model Hamiltonian is given in eq. (51.1), which can be written formally as

$$H = \epsilon g_2 + g g_4 \quad (92)$$

in our notation. This equation is solved using the mean field ansatz. It is convenient to introduce the following functions (FW eq.s 51.10, 51.12a, 51.12b, 51.14)

$$\begin{aligned} \mathcal{G}(x, x') &:= - \langle T[\psi_{k\uparrow} \psi_{k\uparrow}^\dagger] \rangle \quad (\cong \langle a_i a_i^\dagger \rangle_{\nu w}) \\ \mathcal{F}(x, x') &:= - \langle T[\psi_{k\uparrow} \psi_{k\downarrow}] \rangle \quad (\cong \langle a_1 a_2 \rangle_{\nu w}) \\ \mathcal{F}^\dagger(x, x') &:= - \langle T[\psi_{k\downarrow}^\dagger \psi_{k\uparrow}^\dagger] \rangle \quad (\cong \langle a_2^\dagger a_1^\dagger \rangle_{\nu w}) \\ \Delta(x) &:= g \mathcal{F}(x, x). \end{aligned} \quad (93)$$

Δ is the gap-function, which afterwards describes the lowering of energy in the superconducting phase. In our above notation, we had F as propagator, which corresponds to the Green function $\mathcal{G}$ while $\mathcal{F}$ and $\mathcal{F}^\dagger$ have to be identified with the shift $\Delta_0 = uu^\dagger$. This renders $|u|$ to be the gap for a single mode. However, as we saw in the Clifford geometric approach, this parameter is not responsible for the general structure of the vacua in the ν - w -plane, but induces only a shift. Unfortunately the Bogoliubov-Valatin transformation and the BCS theory loose this connection by mixing operators and not properly mixing the states. If we introduce $\xi_k = \epsilon_k - \mu = \hbar^2 k^2 / 2m - \mu$ (FW eq. 37.24 and 51.29) as the energy relative to the chemical potential, one can give the above functions in momentum space as (FW eq.s 51.30a, 51.30b)

$$\begin{aligned} \mathcal{G}(k, \omega) &= \frac{-\hbar(i\hbar\omega + \xi_k)}{\hbar^2\omega^2 + \xi_k^2 + |\Delta|^2} \\ \mathcal{F}(k, \omega) &= \frac{\hbar\Delta^*}{\hbar^2\omega^2 + \xi_k^2 + |\Delta|^2} \end{aligned} \quad (94)$$

where the gap Δ is real in absence of an external field. As a last step, one has to specify these functions as functions of the parameters of the Bogoliubov-Valatin transformation. This yields (FW eq.s 51.31–51.34)

$$\begin{aligned} E_k &= \sqrt{\xi_k^2 + \Delta^2} \\ v_k u_k &= \frac{\Delta}{2E_k} \\ v_k^2 &= 1 - u_k^2 = \frac{1}{2} \left(1 - \frac{\xi_k}{E_k} \right) \\ \mathcal{G}(k, \omega) &= \frac{u_k^2}{i\omega_k - E_k/\hbar} + \frac{v_k^2}{i\omega_k + E_k/\hbar} \\ \mathcal{F}(k, \omega) &= \mathcal{F}^\dagger(k, \omega) \\ &= u_k v_k \left[\frac{1}{i\omega_k - E_k/\hbar} - \frac{1}{i\omega_k + E_k/\hbar} \right]. \end{aligned} \quad (95)$$

Inserting our expressions for the u_k, v_k in terms of the ρ_k (91) we are able to get all of the results as usual. However, this discussion makes it quite clear, that the common treatment of BCS theory involves several peculiar steps. First of all, it was unveiled that the BCS and/or Bogoliubov-Valatin treatment is not able to include the third edge state and deals thus with a restricted theory. Secondly, the BCS and Bogoliubov-Valatin treatment spoils particle number conservation due to the naive concept of vacua involved. A more elaborate treatment which mixes the algebraic states and not the operators can surmount this difficulty. Thirdly, the gap is calculated afterwards by a consistency condition and/or a variational method. The algebraic theory however fixes these parameters uniquely from propagator data and thus ultimatively from the chosen Hamiltonian. This can easily include the thermodynamical behaviour of the theory, if the model Hamiltonian is explicitly known in terms of temperature dependent couplings etc. One ends up with a theory such as

$$H = \alpha(T, B, \dots) \mathbb{1} + \epsilon(T, B, \dots) g_2 + g(T, B, \dots) g_4. \quad (96)$$

Playing with the external parameters $T, B, \dots$ we can move through the vacuum plane, since ν and w become functions of $T, B, \dots$. This explains also, in which way a phase transition can be experimentally obtained. However, the dependence of the coupling constants on these thermodynamical variables can be derived only from a more subtle microscopic theory which is beyond BCS theory. One should also compare the treatment of superconductivity in [42].

4.3 Summary

By embedding CAR algebras and Extended CAR algebras into a generalized Clifford geometric framework it was possible to unveil several new aspects. This fact is based on an intrinsic algebraization of states in Clifford algebras of multivectors. If we have a duality pairing $\langle \cdot, \cdot \rangle$ and an exterior product $\wedge$, we define by

$$\langle a \lrcorner b, c \rangle = \langle b, a^\sim \wedge c \rangle \quad (97)$$

the contraction as the dual product of the wedge w.r.t. the given duality pairing. On the other hand, we might fix a contraction and a wedge, thereby defining a duality pairing. This cannot be achieved if the positivity requirement is considered, since the positive states constitute only a convex affine set, which is i.g. not linear. Furthermore, we can not find a representation of the full algebraic setting in a positive nondegenerate space of appropriate dimension, which is seen from the GNS construction and Clifford algebra theory. However, we saw that ECAR methods are able to describe correctly the vacuum sector, but looses thereby the *constructive* relation between the propagator matrix elements and the vacuum parameterization. This relation is uncovered by introducing Clifford algebras of multivectors or quantum Clifford algebras, which possess possibly an antisymmetric part in the contraction. In general, this introduces $n(n-1)/2$ parameters, which can be used to describe inequivalent vacua. However, since one has already chosen a basis, these propagator matrix elements are basis dependent and thereby observer dependent, so that not all of them describe properties of vacua. We succeeded in our $U(2)$ -model, to give the explicit functional relations of the

vacuum parameters ν, w in terms of the contraction data. As we were able to parameterize all possible propagator we can describe all possible dynamics by five parameters: $\nu, w, \Delta_0, \Delta_1; q$.

We were able to give a complete classification of the vacuum sector of positive states, including a phase transition. We obtained, furthermore, shifts in the gap-equation induced by spin-one and spin-zero eigenvectors. No additional ordering parameter or potential was needed to obtain this transition, which occurs for *purely algebraic reasons*. This ordering parameter is replaced by the Higgs field in elementary particle theory. Moreover, we did not even need to define a specific dynamics but used the propagator as a parameter. This renders the vacuum structure to be universal to any models or choices of Hamiltonians possible within our framework.

Since our method can formally be generalized to QFT in a straightforward manner [15, 13] and even to symplectic Clifford algebras, which are related to CCR algebras and bosons [15], we conclude that symmetry breaking might occur without a Higgs field. The Higgs field of elementary particle physics would therefore *ad hoc* reintroduce the lost connection between contraction data and the vacua.

Our model shows very clearly that phase transitions do not depend on an infinite number of particles or a thermodynamic limit.

The relation between operators and observables has been clarified by explicit calculations, which lead e.g. to an *additive* renormalization of eigenmonomials in the generators. Furthermore, our method is *not* restricted to the positive case. Positivity was only considered for comparing our results with conventional calculations. If we allow indefinite states, we have to move to the geometrical point of view, since the statistical interpretation breaks down. In QCD and non-linear spinor field theories ghost fields occur, e.g. the Faddeev-Popov-fields, which are needed during interaction, but not in the initial and final states, which are subjected to a statistical interpretation. We hope that the geometric concepts, valid in indefinite situations, will be useful in the investigation of this situation.

Further developments of the theory shall include explicitly dynamical models with special Hamiltonians. This will then allow us to calculate the propagator matrix elements and therewith the correct and consistent ground state. This will be interesting in non-linear dynamics which intrinsically specifies all data.

4.4 Epilogue

Our method as presented in this article remains in the area of algebraic developments. However, recent research has unveiled a deeper link of this method to Hopf algebras. Starting from a pair, product and co-product on a space, one can define a convolution product which might be a bigebra or Hopf gebra if additional requirements as a crossing or an antipode are imposed. One finds, that a product on the dual space induces a co-product on the original space. Since we have induced such a duality via our unique vacuum state and such a duality is already employed, the question arises, if we are dealing here silently with a Hopf gebra. One notes, that only with the help of quantum Clifford algebras it is possible to stay with a Hopf gebra structure [23]. If the scalar and co-scalar product are mutually inverse one of another, then only a Clifford convolution algebra can be defined. However, for Grassmann algebras a unique and natural Hopf

algebra structure is easily established. This structure allows one to find deeper arguments what is going on e.g. in re-ordering in QFT [24]. There it was demonstrated that our vacuum is nothing but the co-unit of the involved Hopf algebra and that a process called cliffordization [40] connects the different sets of bases which we have parameterized by v or v, w in our examples. This outcome supports our point that dynamics should select a unique ‘vacuum’ state, i.e. a unique co-unit. Having the power of the Hopf algebra at our disposal, one can turn the usual arguments in QFT upside down and seek for *axioms* which provide models which can be treated as QF theories. In [22] we gave a first account how such axioms might be set up to avoid singularities in QFT by algebraic design. The present work is thus the foundation of such an algebraic axiomatization of quantum physics, which renders various QF theories to be instances of sets of algebraic axioms.

5 Acknowledgements

This paper was inspired by the possibility to play with “Clifford” Ver. 4, a Maple V Release 5 add-on programmed by Prof. Rafal Ablamowicz, Tennessee [2]. His generous help in extending the features, collected in the Cli4plus package, were a valuable help. The support of Mrs. Ursula Wieland is gratefully acknowledged. My gratitude is expressed to Ms. Kirsten Magee for improving my English.

References

- [1] Ablamowicz, R., Lounesto, P., ‘On Clifford algebras of a bilinear form with an antisymmetric part’ in Clifford algebras with Numeric and Symbolic Computations, R. Ablamowicz, P. Lounesto, J.M. Parra (eds.), 167–188, Birkhäuser/Boston 1996.
- [2] Ablamowicz, R., ‘Clifford algebra computations with Maple’ in proceedings of Clifford (Geometric) Algebras, Banff 1995, Alberta Canada, W.E. Baylis ed. Birkhäuser/Boston, 463–501, 1996.
available at: <http://math.tntech.edu/rafal/>
- [3] Ablamowicz, R., Fauser, B., ‘Hecke algebra representations in ideals generated by q -Young Clifford idempotents’ in Clifford Algebras and their Applications in Mathematical Physics, R. Ablamowicz, B. Fauser, Eds., Birkhäuser, Boston, 245–268, 2000, (math.QA/9908062).
- [4] Anderson, P.W., ‘Random-phase approximation in the theory of superconductivity’, *Phys. Rev.* 112, 1900–1916, (1958).
- [5] Araki, H., Woods, E.J., ‘Representations of canonical commutation relations describing a nonrelativistic infinite free bose gas’ *J. Math. Phys.* 4, 637–662, (1963).
Araki, H., Wyss, W., ‘Representation of canonical anticommutation relations’ *Helv. Phys. Acta.* 37, 136–159, (1964).

- [6] Bogoliubov, N.N., ‘On a new method in the theory of superconductivity’, *Nuovo Cim.* 7, 794–805, (1958).
Bogoliubov, N.N., Tolmachov, V.V., Sirkov, D.V., ‘A new method in the theory of superconductivity’, *Fortschritte der Physik* Vol 6, 605–682, (1958).
- [7] Bratelli, O., Robinson, D.W., *Operator algebras and quantum statistical mechanics*, Vols. I,II., Springer, Berlin, 1979, 1981.
- [8] Chevalley, C., *The algebraic theory of spinors*, Columbia University Press, New York, 1954.
- [9] Coxeter, H.S.M., Moser, W.O.J., *Generators and relations for discrete groups*, Springer, Berlin, 1957, [3rd ed.1980].
- [10] Dirac, P.A.M., *Principles of quantum mechanics*, Oxford University Press, Oxford, 1930.
- [11] Fairlie, D.B., Zachos, C.K., ‘Multiparameter associative generalizations of canonical commutation relations and quantized planes’, *Phys. Lett. B* 256(1), 43–49, (1991).
- [12] Fauser, B., ‘Vertex normal ordering as a consequence of nonsymmetric bilinearforms in Clifford algebras’, *J. Math. Phys.* 37(1), 72–83, (1996), (hep-th/9504055).
- [13] Fauser, B., *Clifford-algebraische Formulierung und Regularität der Quantenfeldtheorie*, thesis Tübingen Univ., (1996).
- [14] Fauser, B. ‘Clifford algebraic remark on the Mandelbrot set of two-component number systems’, *Adv. in Appl. Clifford Alg.* 6(1), 1–26, (1996), (hep-th/9507133).
- [15] Fauser, B., Stumpf, H., ‘Positronium as an example of algebraic composite calculations’ in proceedings of the International Conference on the Theory of the Electron, Cuautitlan, Mexico 1995, J. Keller, Z. Oziewicz (eds.) *Advances in Applied Clifford Algebras* 7(S), 399–418, (1997), (hep-th/9510193).
- [16] Fauser, B., ‘Dirac theory from a field theoretic point of view’, in *Clifford algebras and their application in mathematical physics*, IV. Workshop held at Aachen 1996, (V. Dietrich, K. Habetha, G. Jank (eds.)) Kluwer Accad. Publ./Dordrecht, 89–107, 1998.
- [17] Fauser, B., ‘On an easy transition from operator dynamics to generating functionals by Clifford algebras’, *J. Math. Phys.* 39(9), 4928–4947, (1998).
- [18] Fauser, B., ‘On the relation of Clifford-Lipschitz groups to q -symmetric groups’, *Int. Colloquium on Grp. Theor. Meth. in Phys.* 13.–17. in Hobart, Australia July 1998, S.P. Corney, R. Delbourgo, P.D. Jarvis Eds., International Press, Cambridge MA., 413–417, (1999), (q-alg/9807158).

- [19] Fauser, B., ‘Hecke algebra representations within Clifford geometric algebras of multivectors’, *J. Phys. A: Math. Gen.* 32, 1919–1936, (1999).
- [20] Fauser, B., Ablamowicz, R., ‘On the decomposition of Clifford algebras of arbitrary bilinear form’ in *Clifford Algebras and their Applications in Mathematical Physics*, R. Ablamowicz, B. Fauser, Eds., Birkhäuser, Boston, 341–366, 2000, (math.QA/9911180).
- [21] Fauser, B., ‘On the relation of Manin’s quantum plane and quantum Clifford algebras’ *Czech. J. Phys.* 50(1), 1221–1228, (2000), (math.QA/0007137).
- [22] Fauser, B., ‘Quantum Clifford Hopf algebra for quantum field theory’, talk at the Symposium on Quantum Groups, GROUP23, Dubna, August 2000, (hep-th/0011026).
- [23] Fauser, B., Oziewicz, Z., ‘Clifford Hopf algebra for two-dimensional space’, preprint, (math.QA/0011263).
- [24] Fauser, B., ‘On the Hopf algebraic origin of Wick normal-ordering’, *J. Phys. A: Math. Gen.* 34(1), 105–115, (2001), (hep-th/0007032).
- [25] Fetter, A.L., Walecka, J.D., *Quantum theory of many-particle systems*, McGraw-Hill Inc., New York, 1971.
- [26] Fock, V., Iwanenko, D., ‘Über eine mögliche geometrische Deutung der relativistischen Quantentheorie’, *Z. Physik* 54, 789, (1929).
‘Géométrie quantique linéaire et déplacement parallèle’, (presented by Maurice de Broglie), *C.R. Acad. Science (Paris)* 188, 1470, (1929).
Fock, V., ‘Geometrisierung der Diracschen Theorie des Elektrons’, *Z. Physik* 55, 261, (1929).
‘Sur les équations de Dirac dans la théorie de relativité générale’, *C.R. Acad. Sciences (Paris)* 189, 25, (1929).
- [27] Haag, R., ‘On quantum field theories’, *Mat. Fys. Medd. Dann. Vid. Selsk.* 29(12), (1955).
- [28] Haag, R., *Local quantum physics*, Springer, Berlin, 1992.
- [29] Hestenes, D., ‘A unified language for mathematics and physics’ in *Clifford algebras and their applications in mathematical physics*, I. Workshop held at Canterbury, UK, 1985, J.S.R. Chisholm, A.K. Common (eds.), Kluwer, Dordrecht, p.1, 1986.
- [30] Holstein, T., Primakoff, H., ‘Field dependence of the intrinsic domain magnetization of a ferromagnet’, *Phys. Rev.* 58, 1098, (1940).
- [31] Kerschner, R., *Effektive Dynamik zusammengesetzter Quantenfelder*, thesis, Tübingen Univ., 1994, Chap. 4.

- [32] Kerschner, R., ‘On quantum field theories with finitely many degrees of freedom’, unpublished, (hep-th/9505063).
- [33] Lounesto, P., Clifford algebras and spinors, Cambridge University Press, Cambridge, 1997.
- [34] Majid, S., Foundations of quantum group theory, Cambridge University Press, Cambridge, 1995, Chap. 3.
- [35] Murphy, G.J., C^* -algebras and operator theory, Academic Press Inc., San Diego, 1990.
- [36] von Neumann, J., Mathematische Grundlagen der Quantenmechanik, Springer, Berlin, 1932.
- [37] Oziewicz, Z., ‘From Grassmann to Clifford’ in Clifford algebras and their application in mathematical physics, I. Workshop held at Canterbury, UK, J.S.R Chisholm, A.K. Common (eds.), Kluwer, 245–256, 1986.
- [38] Oziewicz, Z., ‘Clifford algebra of multivectors’ in proceedings of the International Conference on the Theory of the Electron, Cuautitlan, Mexico 1995, J. Keller, Z. Oziewicz (eds.) *Advances in Applied Clifford Algebras* 7(S), 1–20, (1997).
- [39] Riesz, M., Clifford numbers and spinors, The Institute of Fluid Dynamics and Applied Mathematics, Lecture Series No. 38, University of Maryland, 1958. Reprinted as facsimile (E.F. Bolinder, P. Lounesto eds.) Kluwer, Dordrecht 1993.
- [40] Rota, G.-C., Stein, J.A., ‘Plethystic Hopf algebras’, *Proc. Natl. Acad. Sci. USA* 91(26), 13062–13066, (1994).
- [41] Streater, R.F., Wightman, A.S., PCT, spin and statistics and all that, Benjamin, New York, 1964.
- [42] Stumpf, H., Borne, Th., Composite particle dynamics in quantum field theory, Vieweg, Braunschweig, 1994.
- [43] Valatin, J.G., ‘Comments on the theory of superconductivity’, *Nuovo. Cim.* VII, 843–857, (1958).
- [44] Wightmann, A.S., ‘Quantum field theories in terms of vacuum expectation values’, *Phys. Rev.* **101**, 860–866, 1956.




Figure 1: Simplex of the positive vacua in the ν - w -plane. The interior of the triangle constitutes of all possible non-degenerate vacua of the $U(2)$ -model. The quasi-free states possessing no higher correlations are described by the parabolas $(\nu, \nu^2 \pm \Delta_i)$. Edge states are pure states and maximally degenerate. Fock, dual-Fock and the new edge states span the simplex. States on the borderline of the simplex are also degenerate, especially the Bogoliubov-Valatin states which connect the dual-Fock and Fock edges. The line connecting the edge-state and a Bogoliubov-Valatin state crosses various parabolas of quasi-free states according to the relevant shifts. This is the path of the bifurcation diagram discussed in the text. States below these parabolas do not have a condensate, while above it a condensate exists. The shifts Δ_i move the parabolas down and up, thereby increasing or decreasing the area of condensate states. In effect, no condensate or only condensate states are possible.